

Fetal Development

4 ½ weeks: heartbeat
6 weeks: brain activity recorded
12 weeks: all major body parts are present
16 weeks: can suck his/her thumb
20 weeks: can feel pain, possibly as early as 12wks
21 weeks: can live outside the womb
24 weeks: hiccupping and smiling
26 weeks: sleep patterns exist
30 weeks: Mom's voice may be recognized
38 weeks: considered full term



Scan to learn more about fetal development or visit
<https://rtl.org/abortion/fetal-development/>




First Trimester
0-13 weeks

Second Trimester
14-26 weeks

Third Trimester
27-40 weeks

Types of Abortion

Suction Abortion- one of the most common methods in the first 12 weeks
RU-486- Chemical abortion used through the 7th week
Dilation & Curettage (D&C)- up to the 12th week
Dilation and Evacuation (D&E)- up to 18 weeks
Dilation and Extraction (D&X or Partial Birth Abortion)- 2nd and 3rd Trimester
Hysterotomy- used the last 3 months of pregnancy (about 28 weeks)
Intracardial Injection- performed at 4 months (about 16 weeks)
Prostaglandin- used during second half of pregnancy (after 20 weeks)
Saline Injection- used during months 4-7 (16-28 weeks)
Heart Attack Abortion- late term abortion



Fetal Development

4 ½ weeks: heartbeat
6 weeks: brain activity recorded
12 weeks: all major body parts are present
16 weeks: can suck his/her thumb
20 weeks: can feel pain, possibly as early as 12 wks
21 weeks: can live outside the womb
24 weeks: hiccupping and smiling
26 weeks: sleep patterns exist
30 weeks: Mom's voice may be recognized
38 weeks: considered full term



Scan to learn more about fetal development or visit
<https://rtl.org/abortion/fetal-development/>



First Trimester
0-13 weeks

Second Trimester
14-26 weeks

Third Trimester
27-40 weeks

Types of Abortion

Suction Abortion- one of the most common methods in the first 12 weeks
RU-486- Chemical abortion used through the 7th week
Dilation & Curettage (D&C)- up to the 12th week
Dilation and Evacuation (D&E)- up to 18 weeks
Dilation and Extraction (D&X or Partial Birth Abortion)- 2nd and 3rd Trimester
Hysterotomy- used the last 3 months of pregnancy (about 28 weeks)
Intracardial Injection- performed at 4 months (about 16 weeks)
Prostaglandin- used during second half of pregnancy (after 20 weeks)
Saline Injection- used during months 4-7 (16-28 weeks)
Heart Attack Abortion- late term abortion




Scan to learn the statistics, methods, and risks of abortion or visit <https://www.prolifefuture.org/abortion-facts/abortion-facts>



Educate yourself on what you are advocating for or against.
<https://www.liveaction.org/news/watched-pro-life-videos-time/>
Viewer discretion advised.



The Aftermath of Abortion



There is help in Michigan!
Scan for resources for pregnancy help in Michigan or visit <https://rtl.org/pregnancy-help/>



**BE EDUCATED AND LET YOUR ACTIONS FIGHT
AGAINST THIS ATROCITY!**

Brought to you by Pro-Life Michigan: <https://prolifemichigan.org/>




Scan to learn the statistics, methods, and risks of abortion or visit <https://www.prolifefuture.org/abortion-facts/abortion-facts>



Educate yourself on what you are advocating for or against.
<https://www.liveaction.org/news/watched-pro-life-videos-time/>
Viewer discretion advised.



The Aftermath of Abortion



There is help in Michigan!
Scan for resources for pregnancy help in Michigan or visit <https://rtl.org/pregnancy-help/>



**BE EDUCATED AND LET YOUR ACTIONS FIGHT
AGAINST THIS ATROCITY!**

Brought to you by Pro-Life Michigan: <https://prolifemichigan.org/>